

9. **Nason GJ, O’Kelly F, Mulvin DW.** Double trouble – ureteric obstruction due to bilateral urothelial carcinoma. *Ir Med J.* 2013; 106(6): 182–183.
10. **Bravo Fernández I, García Cardoso JV, Pérez Ortiz G, et al.** Synchronous bilateral urothelial carcinoma of the upper urinary tract. *Actas Urol Esp.* 2006; 30(3): 335–339.
11. **Fang D, Xiong G, Li X, et al.** Incidence, characteristics, treatment strategies, and oncologic outcomes of synchronous bilateral upper tract urothelial carcinoma in the Chinese population. *Urol Oncol.* 2015 Feb; 33(2): 66.e1–11. Epub 2014 Sep 16.
12. **Látalová B, Grepl M, Čtvrtlík F, et al.** Rychle progredující uroteliální tumor pánvičky levé ledviny s trombózou vena cava inferior. *Ces Urol* 2015; 19(1): 78–82.
13. **Maestroni U, Ferretti S, Dinale F, et al.** A renal cancer with intermediate characteristics between collecting (Bellini) duct carcinoma and urothelial carcinoma: case report and review of the literature. *Tumori.* 2006; 92(6): 545–548.
14. **Pécuchet N, Bigot F, Gachet J, et al.** Triple combination of bevacizumab, gemcitabine and platinum salt in metastatic collecting duct carcinoma. *Ann Oncol.* 2013 Dec; 24(12): 2963–2967. Epub 2013 Nov 4.
15. **Zhu L, Wang Z, Pan C, et al.** Surgical monotherapy may be a suitable therapeutic strategy for advanced collecting (Bellini) duct carcinoma: a case report and literature review. *Exp Ther Med.* 2016; 12(2): 1181–1184.