

16. Pritchett TR, Moreno J, Warner NE, et al. Undiscovered prostatic adenocarcinoma in patients who have undergone radical cystoprostatectomy for transitional cell carcinoma of the bladder. *J Urol* 1988; 139(6): 1214–1216.
17. Epstein JI, Walsh PC, Carmichael M, Brendler CB. Pathologic and clinical findings to predict tumor extent of nonpalpable (stage T1c) prostate cancer. *Jama* 1994; 271(5): 368–374.
18. Epstein JI, Pizov G, Walsh PC. Correlation of pathologic findings with progression after radical retro-pubic prostatectomy. *Cancer* 1993; 71(11): 3582–3593.
19. Epstein JI, Carmichael M, Partin AW, Walsh PC. Is tumor volume an independent predictor of progression following radical prostatectomy? A multivariate analysis of 185 clinical stage B adenocarcinomas of the prostate with 5 years of followup. *J Urol* 1993; 149(6): 1478–1481.
20. Stamey TA, Freiha FS, McNeal JE, et al. Localized prostate cancer. Relationship of tumor volume to clinical significance for treatment of prostate cancer. *Cancer* 1993; 71(3 Suppl): 933–938.
21. Bastian PJ, Mangold LA, Epstein JI, Partin AW. Characteristics of insignificant clinical T1c prostate tumors. A contemporary analysis. *Cancer* 2004; 101(9): 2001–2005.
22. Hekal IA, El-Tabey NA, Nabeeh MA, et al. Validation of Epstein criteria of insignificant prostate cancer in Middle East patients. *Int Urol Nephrol* 2010; 42(3): 667–671.
23. Lee MC, Dong F, Stephenson AJ, et al. The Epstein criteria predict for organ-confined but not insignificant disease and a high likelihood of cure at radical prostatectomy. *Eur Urol* 2010; 58(1): 90–95.
24. Epstein JI, Allsbrook WC, Jr Amin MB, Egevad LL. The 2005 International Society of Urological Pathology (ISUP) Consensus Conference on Gleason Grading of Prostatic Carcinoma. *Am J Surg Pathol* 2005; 29(9): 1228–1242.
25. Epstein JI. An update of the Gleason grading system. *J Urol* 2010; 183(2): 433–440.
26. Chen ME, Johnston D, Reyes AO, et al. A streamlined three-dimensional volume estimation method accurately classifies prostate tumors by volume. *Am J Surg Pathol* 2003; 27(10): 1291–1301.
27. Carvalhal GF, Humphrey PA, Thorson P, et al. Visual estimate of the percentage of carcinoma is an independent predictor of prostate carcinoma recurrence after radical prostatectomy. *Cancer* 2000; 89(6): 1308–1314.
28. Pan J, Xue W, Sha J, et al. Incidental prostate cancer at the time of cystectomy: the incidence and clinicopathological features in Chinese patients. *PLoS One* 2014; 9(4): e94490.
29. Desai A, Wu H, Sun, L, et al. Complete embedding and close step-sectioning of radical prostatectomy specimens both increase detection of extra-prostatic extension, and correlate with increased disease-free survival by stage of prostate cancer patients. *Prostate Cancer Prostatic Dis* 2002; 5(3): 212–218.
30. Fritsche HM, Aziz A, Eder F, et al. Potentially clinically relevant prostate cancer is found more frequently after complete than after partial histopathological processing of radical cystoprostatectomy specimens. *Virchows Arch* 2012; 461(6): 655–661.