

2. Kolon TF, Herndon CD, Baker LA, et al. Evaluation and treatment of cryptorchidism: AUA guideline. *The Journal of urology* 2014; 192(2): 337–345.
3. EAU Guidelines for Paediatric Urology. <http://uroweb.org/guideline/paediatric-urology/>.
4. Šmakal O, Zapletalová J. Polohové anomálie varlat – diagnostika a léčba. *Urol. praxi* 2002; 3(2): 49–51.
5. Hutson JM, Southwell BR, Li R, et al. The Regulation of Testicular Descent and the Effects of Cryptorchidism. *Endocr Rev* 2013; 34(5): 725–752. doi: 10.1210/er.2012–1089.
6. Pettersson A, Richiardi L, Nordenskjöld A, Kaijser M, Akre O. Age at Surgery for Undescended Testis and Risk of Testicular Cancer. *N Engl J Med* 2007; 356(18): 1835–1841.
7. Tasian GE, Copp HL. Diagnostic performance of ultrasound in nonpalpable cryptorchidism: a systematic review and meta-analysis. *Pediatrics* 2011; 127(1): 119–128.
8. Elder JS. Ultrasonography is unnecessary in evaluating boys with a nonpalpable testis. *Pediatrics* 2002; 110: 748–751.
9. Adesanya OA, Ademuyiwa AO, Evbuomwan O, Adeyomoye AA, Bode CO. Preoperative localization of undescended testes in children: Comparison of clinical examination and ultrasonography. *J Pediatr Urol* 2014; 10(2): 237–240.
10. Nijs SM, Eijbouts SW, Madern GC, et al. Nonpalpable testes: is there a relationship between ultrasonographic and operative findings? *Pediatr Radiol* 2007; 37: 374–379.
11. Berger C, Haid B, Becker T, et al. Nonpalpable testes: Ultrasound and contralateral testicular hypertrophy predict the surgical access, avoiding unnecessary laparoscopy. *J Pediatr Urol* 2017; 24: 3056–63.
12. Bae KH, Park JS, Jung HJ, Shin HS. Inguinal approach for the management of unilateral non-palpable testis: is diagnostic laparoscopy necessary? *Journal of pediatric urology* 2014; 10(2): 233–236.
13. Budianto IR, Tan HL, Kinoshita Y, et al. Role of laparoscopy and ultrasound in the management of „impalpable testis“ in children. *Asian journal of surgery* 2014; 37(4): 200–204.
14. Snodgrass W, Chen K, Harrison C. Initial scrotal incision for unilateral nonpalpable testis. *J Urol* 2004; 172: 1742e5.
15. Shepard CL, Kraft KH. The Nonpalpable Testis: A Narrative Review. *J Urol* 2017; 198(6): 1410–1417.
16. Akihiro I, Kikuchi K, Ougushi K, et al. Surgical exploration for impalpable testis: Which should be first, inguinal exploration or laparoscopic abdominal exploration. *Journal of Pediatric Surgery* 2018; 53(9): 1766–1769.
17. Drlík M, Kočvara R. Nehmatné varle u dítěte: začít diagnostickou laparoskopií nebo otevřenou revizí třísla? *Ces Urol* 2011; 15(2): 101–107.